

## Characterization and Identification of A Sludge-Associated Bacterial Isolate

**Abiodun O. Adibi**<sup>1</sup> and **Derrell McPherson**, Department of Biological Sciences, Hampton University, Hampton, VA 23668, and  
**Minna Laine**, Finnish Environment Agency Laboratory,  
Hakuninmaantie 4-6, FIN-00430 Helsinki, Finland.

### ABSTRACT

During a metal speciation study, an unusually high retention of lead was observed when lead solution was percolated through a column packed with sludge compost obtained from Hampton Roads Sanitation department. A bacterium was isolated from the sludge compost and identified as *Bacillus sphaericus* using electron microscopy, whole cell fatty acid analysis (Midi System) and Biolog GP Microplate. The isolate grows in broth and agar media containing up to 800  $\mu\text{M}$  lead. Lead accumulation study using atomic absorption spectrophotometer indicates that the isolate adsorbs lead. Lead adsorption is pH dependent. The isolate contains a plasmid of approximately 40 – 50 kbp that might be involved in resistance to lead. Studies are in progress to characterize this plasmid. The bacterial isolate has the potential of being used in bioremediation of heavy metal contaminated water and could be involved in localized accumulation of lead in gardens if heavy metal containing sludge compost is used as fertilizer.

### INTRODUCTION

Composted wastewater sludge serves as sources of nutrients when applied to lawns and gardens. This method of disposal of wastewater sludge is currently preferred to incineration and landfilling. There is variation in bacterial composition of sludge. Some of the most frequently found bacteria in sewage are in the genera *Beggiatoa*, *Achromobacter*, *Sphaerotilus*, *Flavobacterium*, *Pseudomonas*, *Escherichia*, *Enterobacter* and *Zooglea*. However, this bacterial composition changes during sewage treatment and when sludges obtained from sewage treatment plants are used in compost production. Temperature generated during composting results in the displacement of mesophilic bacteria by thermophilic bacteria. The latter group includes species of *Clostridium*, *Thermoactinomyces* and *Bacillus*. Studies have shown that wastewater sludge contain high concentrations of cations. A study of the residual metal contents of sludge from Hampton Roads in Virginia revealed the presence of copper, iron, manganese, lead and zinc. Thus, sludge-composts are potential sources of bacterial and heavy metals contamination of water and soil.

The roles of microorganisms in heavy metal uptake are subjects of many studies. These studies include the uptake of Mn, Sr, Zn, Cu and Cd by *Saccharomyces cerevisiae* (Avery and Towbin, 1993); accumulation of cesium from soil by two

---

1 Corresponding author: Phone # (757) 727-5017, E-mail address: facadibi@hamptonu.edu

*Rhodococcus* sp. (Tomioka et al., 1994); accumulation of nickel and zinc by filamentous bacterium in the genus *Thiothrix* (Shuttleworth and Unz, 1993); accumulation of copper by yeasts (Junghans and Straube, 1991). Studies conducted by Mullen et al. (1989) showed that *Bacillus subtilis*, *B. cereus*, *Escherichia coli* and *Pseudomonas aeruginosa* were able to remove one or more of a variety of cations from solutions. Biosorption of metals by microorganisms occurs by different mechanisms and is affected by different factors. Metabolism-independent biosorption in some microorganisms occurs on the cell wall and capsule (Shuttleworth and Unz, 1993; Dyer et al., 1994; Avery and Tobin, 1993; Ortega-Calva et al., 1994; Gadd, 1990; Scott and Palmer, 1988). According to Silver and Walderhaug (1992), genes located on bacterial chromosomes are responsible for bacterial accumulation of several cations and anions. The group also showed that resistances to some heavy metals are plasmid-mediated. Presence of other ions may also influence metal uptake by some microorganisms. As was reported by Macaskie et al. (1994), accumulation of heavy metals by *Citrobacter* sp. was mediated by the presence of phosphate in the periplasmic space. Drastic increase in mercuric chloride uptake by genetically engineered *Escherichia coli* was observed in the presence of Na (Selifonova and Barkay, 1994). These show that bacteria introduced into the soil could promote localized concentration of heavy metals, which could have serious consequences on agriculture. Also that bacteria have great potential in bioremediation of heavy metal contaminated environments.

During the course of metal speciation studies conducted using a column packed with sludge compost, an unusually high retention of lead by the column was observed when a lead solution was percolated through the column. Bacteriological examination of the compost revealed the presence of a bacterium, which was isolated and characterized. In this report, we describe the characterization and identification of a bacterial isolate from composted wastewater sludge from Hampton Roads in Virginia.

#### MATERIALS AND METHODS

**Isolation and Growth of Bacterium.** Composted sludge was sifted to remove large organic debris. This was suspended in sterile, phosphate buffered saline, pH 7.2 (PBS) and mixed thoroughly. The suspension obtained was used as inoculum for tryptic soy broth, which was incubated overnight at room temperature. The broth culture was streaked on tryptic soy agar plates and incubated. Bacterial pure culture obtained was stored on tryptic soy agar slant and at -70 C in glycerol (20% v/v in tryptic soy broth).

**Bacterial Plasmid.** Tryptic soy broth containing 400  $\mu$ M lead nitrate was inoculated with a single colony of the isolate and incubated at 30 C. in an orbital shaker incubator. Plasmids were extracted using the rapid alkaline lysis method as described by Birnboim and Doly (1979) with slight modification. Plasmid was digested with restriction endonucleases and analyzed on a 0.6% agarose gel prepared in Tris Acetate EDTA buffer. Gel was stained in ethidium bromide (0.5 mg/L), destained and bands were visualized using the Bio-Rad Gel Doc 1000.

**Physical and Biochemical Characterization.** The following tests were performed on the isolate using standard bacteriological procedures: Gram's stain, growth in different concentrations of NaCl, growth at different pH and temperatures, starch hydrolysis, DNA hydrolysis, citrate utilization, catalase production, indole production, methyl red (MR) and Voges Proskauer (VP).

**Electron Microscopy.** Transmission electron microscopy was done on 24 and 48 hr cultures of the isolate. The isolate was fixed in 2% osmium tetroxide in 0.2M cacodylate buffer, dehydrated in a series of increasing concentration of ethanol, embedded in epon and sectioned using an LKB Ultratome III 8800. Electron microscopy was done using the JEOL 100CX II Transmission Electron Microscope.

**Whole Cell Fatty Acid Analysis.** Fatty acid analysis was performed using the MIDI system. Bacterial isolate was inoculated on Tryptic Soy Broth Agar (TSBA) in petri dishes and incubated at 29° C. Approximately 35mg of the cells were harvested into a precleaned, screw-capped tube. The cell pellet was treated as specified by the manufacture in order to extract cell-associated fatty acids. The extracted fatty acids were resolved using a HP 5890 gas chromatograph and Sherlock Pattern Recognition Software.

**Biolog GP Microplate Analysis.** The Biolog Gram Positive Microplate method using 95 biochemical tests was done according to the manufacturer's direction. The GP Microplate determines the ability of bacterial isolates to oxidize 95 different carbon sources (Figure 1). The isolate was grown in Biolog Universal Medium containing glucose. Bacterial suspension was made in physiological saline and 150 µL of the suspension was inoculated into microtiter plate wells containing different carbon sources and also tetrazolium dye. The microtiter plate was incubated at 30° C for 24 hrs. When the carbon source is utilized there is a blue color formation resulting from reduction of tetrazolium dye incorporated into the medium. Negative wells remain colorless. The result yields a metabolic fingerprint of the inoculated organism

which is entered into Biolog's Microlog computer Identification of isolate is by computerized-comparison of the results with metabolic pattern of library species kept in database.

**Lead Accumulation Studies.** SB-1 was inoculated into LB broth and incubated overnight at 26° C in a shaker incubator. The culture was centrifuged, the pellet washed in sterile deionized water and suspended in water to a known absorbance. This suspension was used as inoculum for screw-capped tubes containing equal volume of physiological saline supplemented with varying concentrations of the metal under study. The tubes were incubated at 26° C in a shaker incubator and samples were retrieved at set intervals of time. The retrieved samples were centrifuged. The supernatants obtained were saved and the pellets washed with distilled water and digested in 1N HNO<sub>3</sub> as described by Shuttleworth and Unz (1993). The metal contents of the digested pellets and the corresponding supernatants were determined using a Perkin-Elmer 4000AAS and the appropriate Intensitron lamp.

## RESULTS AND DISCUSSION

The results of several Gram's staining procedure showed that the bacterial isolate was Gram negative rod. Contrarily, transmission electron microscopy result shown in Figure 2 indicates that the isolate is a sporulating, long rod with round, subterminal endospores. These characteristics are associated with Gram positive bacteria. Old cultures of Gram positive bacteria and Gram positive bacteria with thin peptidoglycan will, in some cases, give a Gram negative result. Electron microscopy shows that the bacterial isolate (SB-1) has thin cell wall. The isolate exhibited very good growth at pH 6.0 – pH 9.0 but very little growth at pH 4.0. Copious growth was evident at up to 4.0% NaCl but very little growth at higher concentration of NaCl. The isolate is

A1	Water	A2	$\alpha$ -cyclodextrin	A3	$\beta$ -cyclodextrin	A4	levirin	A5	Glycogen	A6	Inulin	A7	Mannan	A8	Tween 40	A9	Tween 80	A10	N-acetyl-D-Glucosamine	A11	N-acetyl-B-D-Minosamine	A12	amygdalin
B1	L-Arabinose	B2	D-Arabinofuranose	B3	Arabinofuranose	B4	D-Cellobiose	B5	D-Fructose	B6	L-Fucose	B7	D-Galactose	B8	D-Galacturonic acid	B9	Gentianose	B10	D-Gluconic acid	B11	A-D-Glucose	B12	M-Inositol
C1	A-D-Lactose	C2	Lactulose	C3	Maltose	C4	Maltotriose	C5	D-Mannitol	C6	D-Mannose	C7	D-Maltotriose	C8	D-Melibiose	C9	A-methyl-D-Galactoside	C10	B-methyl-D-Galactoside	C11	3-methyl glucose	C12	A-methyl-D-galactoside
D1	B-methyl-D-Glucoside	D2	Lactulose	D3	Maltose	D4	Maltotriose	D5	D-Raffinose	D6	L-Rhamnose	D7	D-Ribose	D8	Salicin	D9	Sedoheptulose	D10	D-Sorbitol	D11	Stachyose	D12	Sucrose
E1	D-Tangitose	E2	D-Trehalose	E3	Turanose	E4	Xylitol	E5	D-Xylose	E6	Acetic acid	E7	A-Hydroxybutyric acid	E8	B-Hydroxybutyric acid	E9	G-Hydroxybutyric acid	E10	P-Hydroxyphenyl acetic acid	E11	A-keto-Glutaric acid	E12	A-keto-Valeric acid
F1	Lactamide	F2	D-lactic acid Methyl ester	F3	L-Lactic acid	F4	Malic acid	F5	L-Malic acid	F6	Methyl Pyruvate	F7	Methyl Succinate	F8	Propionic Acid	F9	Pyruvic Acid	F10	Succinamic Acid	F11	Succinic Acid	F12	N-acetyl-L-Glutamic acid
G1	Alaninamide	G2	D-Alanine	G3	L-Alanine	G4	L-Alanyl Glycine	G5	L-Asparagine	G6	L-Glutamic Acid	G7	Glycyl-L-Glutamic Acid	G8	L-Pyroglutamic Acid	G9	L-Serine	G10	Purrescine	G11	2,3-Butanediol	G12	Glycerol
H1	Adenosine	H2	2-Deoxy adenosine	H3	Inosine	H4	Thymidine	H5	Uridine	H6	Adenosine-5'-Monophosphate	H7	Thymidine-5'-Monophosphate	H8	Uridine-5'-Monophosphate	H9	Fructose-6-Phosphate	H10	Glucose-1-Phosphate	H11	Glucose-6-Phosphate	H12	D,L-A-Glycerol Phosphate

FIGURE 1. BIOLOG GP Microplate substrates used in SB-1 bacterial identification.



FIGURE 2. Electron micrograph of SB-1 bacterial isolate showing subterminally located endospores in the bacilli.

amylase, oxidase and VP negative but catalase, Dnase and MR positive. Whole cell fatty acid analysis profile of the isolate is shown in Table 1. The result indicates that the isolate is Gram positive because it contains anteiso and iso fatty acids, which are found only in Gram positive bacteria. In addition, the bacterium was identified as *Bacillus sphaericus* based on whole cell fatty acid analysis. This identification was supported by the Biolog GP Microplate results. The similarity index of the Biolog GP Microplate identification was 0.892. A similarity index of at least 0.5 for cultures incubated at 30° C is acceptable for species identification.

The isolate contains a large plasmid that is approximately 40 - 50 kbp. The plasmid has EcoR I, Hind III and Hae III sites but not BamH I Sal I and Sma I sites. Although the function of this plasmid is yet to be determined, the plasmid is believed to be associated with the survival of the isolate in lead and probably other heavy metals. Plasmid preparation from culture grown in nutrient broth without lead resulted in very little yield. On the other hand, culture of the isolate in medium containing up to 40  $\mu\text{M}$  lead resulted in a good yield of plasmid. According to Silver and Ji (1994) and Collard et al. (1994), resistances to heavy metal in some Gram negative bacteria reside in large, naturally occurring plasmids. Studies have begun to characterize the plasmid.

Initial studies indicated that the isolate is capable of growing in culture medium containing up to 600  $\mu\text{M}$  lead, this indicates resistance to lead because lead at very small concentrations is detrimental to bacterial growth. In their studies of bacterial communities in heavy metal contaminated soils, Roane and Kellogg (1996) isolated many lead-resistant genera including *Bacillus*. The mechanism of resistance to the heavy metals was not discussed. The isolate also accumulates lead from solution (Figure 3). Lead accumulation increases with increase in the cell concentration and was pH dependent. Lead accumulation was highest at pH 7.3 and pH 8.0 than at pH 5.0. This finding is supported by Tomioka et al (1994) working with *Rhodococcus* species. They showed that maximum absorption of cesium occurred at pH 8.5. The mode of

TABLE 1. Fatty acid analysis and identification of SB-1 bacterial isolate.

Sherlock Version: 1.06						
ID: 17651 AA-SB-960712						
Bottle:27 SAMPLE (AEROBE)						
RT	Area	Ar/Ht	Respon	ECL	Name	%
1.667	252780000	0.026	----	7.013	SOLVENT PEAK	---
2.828	102	0.027	----	9.543	-----	---
6.367	7974	0.035	1.012	13.618	14:0 ISO	8.62
6.858	768	0.035	0.997	14.000	14:0	0.82
7.769	27366	0.039	0.974	14.621	15:0 ISO	28.46
7.901	13374	0.039	0.970	14.711	15:0 ANTEISO	13.86
8.325	648	0.039	0.961	15.000	15:0	0.67
8.943	11766	0.041	0.949	15.386	16:1 w7c alcohol	11.93
9.326	25146	0.040	0.942	15.625	16:0 ISO	25.30
9.538	3348	0.040	0.938	15.758	16:1 wlle	3.36
9.926	1692	0.044	0.932	16.000	16:0	1.68
10.729	756	0.041	0.920	16.478	Sum In Feature 5	0.74
10.983	1626	0.043	0.917	16.629	17:0 ISO	1.59
11.141	3036	0.044	0.915	16.723	17:0 ANTEISO	2.97
19.600	10008	0.078	----	21.652	-----	---
*****	756	----	----	----	Summed feature 5	.74
Solvent Ar	Total Area	Named Area	% Named	Total Amnt	Nbr Ref	
252780000	79602	97500	99.90	93613	9	

TSBA [Rev 3.90] <i>Bacillus</i>	0.410
<i>B. sphaericus</i>	0.410
<i>B. s.</i> GC subgroup IV*	0.410
<i>B. s.</i> GC subgroup III	0.376

CLIN [Rev 3,90] \* NO MATCH \*

lead biosorption is most likely by adsorption to the cell wall because the lead accumulation studies were conducted using cells suspended in physiological saline. Several studies including that of Strandberg et al., (1981) have shown that bacteria and other microorganisms with cell wall adsorb heavy metals to their cell wall. In a review article, Gadd (1988) indicated that biosorption of heavy metal by *Bacillus subtilis* occurs on the carboxyl group of glutamic acid component of the peptidoglycan. Nakajima and Sakaguchi (1986) showed that *Bacillus subtilis* exhibited extremely high absorption of heavy metals, including lead, copper, uranium, cobalt and nickel. Gram negative bacteria have been the foci of studies for bacterial bioremediation abilities. This study shows that non-pathogenic *Bacillus* species may be important in bioremediation of heavy metal contaminated soil and aquatic environments.




Figure 3. Lead concentrations in SB-1 pellets from solutions containing various lead concentrations (◆, ■, ▲, X and \* represent 10, 20, 40, 80 and 100 mM lead, respectively).

#### ACKNOWLEDGEMENTS

Isolation of SB-1, biochemical and physical characterization, fatty acid analysis and lead accumulation studies were conducted by A. Adibi and Derrell McPherson. Derrell McPherson is currently pursuing a Ph.D. program in Microbiology at Virginia Tech. Minna Laine identified the isolate using the Biology GP method. Thanks are extended to Dr. Isai Urasa and Nixon Nwebi for their assistance with atomic absorption spectrophotometer; also to Miss Betty Penn for her assistance in the preparation of the manuscript.

#### REFERENCES

- Avery, S.V. and Tobin, J.M. 1993. Mechanism of adsorption of hard and soft metal ions to *Saccharomyces cerevisiae* and influence of hard and soft anions. *Appl. Environ. Microbiol.* 59:2851-2856.
- Birnboim, H.C. and Doly, J. 1979. A rapid alkaline procedure for screening recombinant plasmid DNA. *Nucleic Acids Research* 7:1513-1525.
- Collard, J.M., Corbisier, P., Diels, L., Dong, Q., Jeanthon, C., Mergeay, M., Taghavi, S., van der-Lelie, D., Wimotte, A. and Wuertz, S. 1994. Plasmids for heavy metal resistance in *Alcaligenes eutrophus* CH34: mechanisms and applications. *FEMS Microbiol. Rev.* 14:405 - 414.
- Dyer, B.D., Krumbein, W.E. and Mossman, D.J. 1994. Accumulation of gold in the sheath of *Plectonema terebrans* (filamentous, marine cyanobacteria). *Geomicrobiol. J.* 12:91-98.
- Gadd, G.M. 1990. Heavy metal accumulation by bacteria and other microorganisms. *Experientia* 46:834-840.

- Hunt, S. 1986. Diversity of biopolymer structure and its potential for ion binding applications. *In* Immobilization of Ions by Bio-sorption. Edited by H. Eccles and S. Hunt. Ellis Harwood Limited, Chichester.
- Junghans, K. and Straube, G. 1991. Biosorption of copper by yeasts. *Biol. Met.* 4:233-237.
- Macaskie, L.E., Bonthrone, K.M. and Rouch, U.K. 1994. Phosphatase-mediated heavy metal accumulation by a *Citrobacter* sp. and related enterobacteria. *FEMS Microbiol. Lett.* 121:141-146.
- Mullen, M.D., Wolf, D.C., Ferris, F.G., T.J., Flemming, C.A. and Bailey, G.W. 1989. Bacterial sorption of heavy metals. *Appl. Environ. Microbiol.* 55:3143-3149.
- Nakajima, A. and Sakaguchi, T. 1986. Selective accumulation of heavy metals by microorganisms. *Appl. Microbiol. Biotechnol.* 24:59-64.
- Niemi, R.M., Niemela, S.I., Bamford, D.H., Hantula, J., Hyvarinen, T., Forsten, T. and Raateland, A. 1993. Presumptive fecal streptococci in environmental samples characterized by one-dimensional sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Appl. Environ. Microbiol.* 59:2190-2196.
- Rani, D.B. and Mahadevan, A. 1993. Patterns of heavy metal resistance in marine *Pseudomonas* MR1. *Indian J. Expt. Biol.* 31:682-686.
- Roane, T.M. and Kellogg, S.T. 1996. Characterization of bacterial communities in heavy metal contaminated soils. *Can. J. Microbiol.* 42:593-603.
- Scott, J.A. and Palmer, S.J. 1988. Cadmium bio-sorption by bacterial exopolysaccharide. *Biotech. Letters* 10:21-24.
- Selifonova, O.V. and Barkay, T. 1994. Role of  $\text{Na}^+$  in transport of  $\text{Hg}^{2+}$  and induction of the Tn21 mer operon. *Appl. Environ. Microbiol.* 60:3503-3507.
- Shuttleworth, K.L. and Unz, R.F. 1993. Sorption of heavy metals to the filamentous bacterium *Thiothrix* strain A1. *Appl. Environ. Microbiol.* 59:1274-1282.
- Silver, S. 1994. Newer systems for bacterial resistances to toxic heavy metals. *Environ. Health Perspect.* 102 Suppl 3:107-113.
- Silver, S. and Walderhaug, M. 1992. Gene regulation of plasmid- and chromosome-determined inorganic transport in bacteria. *Microbiol. Rev.* 56:195-228.
- Tomioka, N., Uchiyama, H. and Yagi, O. 1994. Cesium accumulation and growth characteristics of *Rhodococcus erythropolis* CS98 and *Rhodococcus* sp. Strain SC402. *Appl. Environ. Microbiol.* 60:2227-2231.